

# ATTRA Identification Sheet

# Apple Diseases



NATIONAL CENTER  
FOR APPROPRIATE  
TECHNOLOGY

There are several key diseases that affect apple production. Use this handy guide to help identify some diseases and to learn more about causes and solutions. For more specific information about these diseases, or to learn about apple diseases not addressed here, see **ATTRA's publication *Apples: Organic Production Guide***.

## — Symptoms —

### Apple Scab

- Apple scab, caused by the fungus *Venturia inaequalis*, causes grayish-brown spots on leaves and hard, rough ("scabby") lesions on fruit. As the lesions progress, they often crack, opening the fruit to other problems.
- Rain, duration of leaf wetness, and temperature determine infection periods.
- If initial infections are not controlled, they will lead to secondary infections later in the season.
- Secondary infections begin when summer spores (conidia) develop in lesions on leaf and bud tissues, to be released during wet periods and disseminated throughout the tree.
- Secondary infections blemish and deform the apples and will also weaken the tree.

### Fire Blight

- Fire blight is caused by the bacteria *Erwinia amylovora*.
- Warm, wet conditions foster the bacteria's reproduction and spread within and among trees. Fire blight will be a problem only in years when weather is conducive to its spread.
- Affected branches wither and turn black or brownish black, as if scorched.
- Most infected branch tips wilt rapidly, taking on the characteristic shape of a "shepherd's crook."
- Under the bark, the bacteria form a canker where they will overwinter.

### Cedar Apple Rust

- The disease produces smallish, round, and yellow to rusty-orange spots on the top and bottom surfaces of apple leaves.
- Susceptible varieties, especially many yellow-fruited varieties, may be defoliated in a year conducive to infection.
- In order to complete its life cycle, the fungus that causes this disease must spend part of its life on eastern red cedar. Orange gelatinous "horns" on the galls on the cedar are actually the fruiting stage of the fungus. The "horns" release the spores that infect the apple trees.

### The Summer Rots

- Characterized by sunken, soft, rotten areas, which sometime consume large parts of the apple.
- Bitter rot typically exhibits a cone-shaped infection, with the base of the cone on the fruit surface.
- The spores of the causal organism of bitter rot sometimes appear in concentric circles in the roundish lesion and, in advanced stages, turn a salmon pink.
- Black rot and white rot are related and similar, but black rot has a leaf phase, frog-eye leaf spot, that leaves black-margined, tan spots on leaves.
- Both black and white rot overwinter on rough patches of bark or in branch crotches, sometimes causing slightly sunken branch cankers. White rot cankers sometimes develop a thin, papery bark.
- Black rot, bitter rot, and white rot are more pervasive in the Southeast than elsewhere, but can become problems elsewhere if the growing season is conducive.

### Sooty Blotch and Fly Speck

- Sooty blotch causes irregular, soot-smear areas on the apple skin.
- Fly speck is characterized by many small, black spots the size of the dot on an "i."
- These two fungal diseases are almost always found together.
- The effects of both diseases are almost purely cosmetic but can render the fruit unsalable.
- These diseases are less apparent on darker fruit, whereas yellow fruit seems to emphasize the problem.



Photo: NCAT



TJ Smith, Washington State University



Paul Bachi, University of Kentucky



Photo: NCAT



Photo: NCAT

# Key Low-Spray and Organic Solutions



## Apple Scab

The use of scab-resistant varieties is the best long-term strategy for organic growers. The number of primary and secondary infections in a year depends on the amount of rain. The warmer the weather, the more quickly conidia development follows primary infection. If the grower is relying on protective-type fungicides, including all organically acceptable fungicides, then trees should be treated whenever there is a chance of primary infection. Apple scab can be controlled on susceptible varieties by timely sprays with fungicides. For the organic apple grower, there are three commonly used materials: sulfur, lime-sulfur, and Bordeaux mixture. Potassium bicarbonate (the trade name is Armicarb) and potassium phosphonate (Resistim), in combination with sulphur, have been shown to be very effective at controlling the fungus. Trees must be sprayed or dusted diligently before, during, or immediately after a rain, from the time of bud break until all the spores are discharged. Because the scab fungus overwinters on fallen apple leaves, growers can largely eliminate the primary scab inoculum and control the disease by raking and destroying the fallen leaves. Neem has also demonstrated some efficacy in managing scab.

## Fire Blight

As of 2014, antibiotics are no longer allowable in organic apple production. Organic growers' first line of defense should be resistant varieties. Next, aim for low-to-moderate tree vigor by avoiding overfertilization and severe pruning. Copper-containing bactericides provide some defense when sprayed at green-tip stage. In the West, the "biological antagonists" Bloomtime Biological<sup>TM</sup> (*Pantoea agglomerans*) and Blossom Protect<sup>TM</sup> (*Aureobasidium pullulans*) sprayed in sequence when the flowers are open provides good control. In the East, these sprays are not as effective (greater disease pressure), but will help in combination with cultural control efforts. During dormancy, fire blight infected wood should be cut out 6-8 inches below the last visible damage and destroyed or removed from the orchard.

## Cedar Apple Rust

Removal of eastern red cedar around the immediate orchard vicinity can reduce the inoculum reaching the apple foliage. In addition, there are many rust-resistant apple varieties. Many fungicides are effective against rust, including the sulfur-and-copper compounds, which are approved for organic production as a last resort in an organic pest-management plan.


## The Summer Rots

For control in organic orchards, growers should emphasize cultural techniques for suppressing the causal organisms of these rots. Such techniques include pruning out diseased wood, removing fruit mummies, pruning for light penetration and air circulation, and avoiding poor sites. Piles of prunings can be a source of infection, so remove prunings from the orchard or burn. Though there is little actual varietal resistance, very early and very late-maturing varieties escape the worst infection periods. A number of biological controls have demonstrated effectiveness against post-harvest rot organisms and are labeled for fruit-rot organisms in general. For post-harvest fruit rots, cultural controls are best combined with a biological antagonist such as *Bacillus subtilis* or *Trichoderma viridi*. Harvesting and storage sanitation practices will help prevent these rots.

## Sooty Blotch and Fly Speck

Planting early-maturing, dark-red apple cultivars can reduce the damage caused by sooty blotch and flyspeck. For the organic orchardist, pruning to maintain adequate air flow through the canopy is very important. Potassium bicarbonate (KHCO<sub>3</sub>) is a new, moderately effective fungicide for control of sooty blotch and fly speck. An amino acid combination of dl-methionine and riboflavin, freshly mixed and supplemented with sodium dodecyl sulfate and a trace amount of copper sulfate, controlled the complex as well or better than the standard sulfur spray. A post-harvest soak for five minutes in a 5 to 10% bleach solution in water has shown effectiveness for sooty blotch control.

Visit [www.attra.ncat.org](http://www.attra.ncat.org) to see ATTRA's *Apples: Organic Production Guide* for more information.



**ATTRA**  
SUSTAINABLE AGRICULTURE

ATTRA was developed and is managed by the National Center for Appropriate Technology (NCAT). The project is funded through a cooperative agreement with the U.S. Department of Agriculture's Rural Business-Cooperative Service.

For more information on a wide range of topics related to sustainable agriculture, visit the ATTRA website at [www.attra.ncat.org](http://www.attra.ncat.org). You'll find hundreds of publications, webinars and videos, news, databases and other tools, and more!