



Integrated Parasite Management: Train the Trainer Project

Grazing to Control Parasites

The Barber Pole Worm has threatened sheep production in the eastern United States and has the potential to impact Intermountain West production as well. In fact, it already has.

Ultimately, combatting the Barber Pole Worm on irrigated pastures with dewormers alone is a losing battle. You will not win because there are only three classes of sheep dewormers and trillions of parasites. The clincher? Time is on their side.

Dewormers have a definite place in a parasite-control program, but how much better would things be if we could drastically reduce the need to use them? Not only will we be saving dewormer, money, and time, we will be improving the soil environment for microbes and arthropods, of which the dung beetle is the most evident. All of these little creatures of the soil do our work for us, improving organic matter, cycling nutrients (see the ATTRA publication *Nutrient Cycling in Pastures* <https://attra.ncat.org/attra-pub/summaries/summary.php?pub=240>) and saving moisture in the soil. That translates into more grass, healthier animals, noticeably less costs, and more profits. But to realize those benefits, we need a plan.

There are two long-term, holistic approaches to reining in the Barber Pole Worm: grazing management and genetic selection. Perhaps the one with the most immediate returns is strategic grazing. We know that the environment plays a huge role (usually 80%) in how a gene is expressed in an animal. For instance, a particular sheep's ability to withstand parasites depends upon her genetic make-up and the environment in which she grazes. If we can create a grazing environment that has fewer Barber Pole Worm larvae to ingest, we and our sheep are miles ahead. Here is the plan.

It has only three rules:

1. An absolute minimum of 35 days of pasture rest. Forty is better.
2. An absolute minimum of 6-inch to 8-inch paddock residual.
3. An absolute maximum of four days in any paddock. One is best.

These rules attack the Barber Pole Worm's life cycle on three fronts: survival, tactical position, and ingestion.

Survival: Paddock Rest Periods of 40 Days

- An adult worm lives about six to eight months inside the sheep abomasum (or stomach), producing several thousand eggs per day. For a quick recap of the Barber Pole Life Cycle, see the ATTRA narrated Power Point, *Don't Let the Barber Pole Worm Devastate Your Flock*. <https://attra.ncat.org/multimedia/ppt>

- *Wintertime*: Unfortunately, we do not start with a clean slate each spring. How do Barber Pole Worms survive our cold Intermountain West winters?


— Inside: L4 larvae (one of the last larval stages) hibernate, or go into hypobiosis, in the folds of the abomasum during the winter, awakening in time for spring lambing and pasture. These larvae then become adults and the egg-shedding (in manure) starts in earnest. Ivermectins (Ivomec®, Eprinex®, Dectomax®) and Moxydectin (Cydectin®) are the most reliable dewormers for killing L4 larvae. Should we deworm all sheep when we come off of fall or winter pasture? Remember the concept of *refugia*. (See the ATTRA tipsheet, *Why FAMACHA® Score?*)

No. Instead, use FAMACHA and only deworm those that score 3, 4, and 5 on the FAMACHA scale.

- Related ATTRA Publication:
 - [**Building Healthy Pasture Soils**](#)

Larval Survival

Survival of barber's pole worm infective larvae on pasture at various daily maximum temperatures and 60% relative humidity



Australian research indicates that 90% of infective larvae die within 40 days of emergence when maximum daily highs reach 35 degrees C or 95 degrees F.

— Outside: Some L3 larvae survive months of frigid temperatures, especially under snow.

- *Summertime*: Research in Australia has discovered that 90% of Barber Pole L3 infective larvae die within 40 days after emerging from eggs when daytime highs are above 90° F or 35° C. (WormBoss, 2018)
- In the summer, when we rest our pasture 40 days or more, we can avoid the peak of Barber Pole larvae populations. It's that easy!
 - A 40-day rest period also sets the stage for optimal plant-nutrient cycling, especially if you graze half and trample half of the forage. This reduces the need for fertilizing, especially nitrogen. (See the ATTRA publication *Nutrient Cycling in Pastures*.)
- Conversely, setting grazing rest at 20 to 25 days (a common practice), syncs grazing periods with peaking parasite populations. Let's avoid that.
- Another effective way to curb parasites is to hay a pasture, which desiccates the parasite larvae. Then, after an adequate rest period, graze that pasture.

Tactical Position: Leave Behind Six to Eight Inches of Grass


- The vast majority of Barber Pole Worm L3 larvae only crawl three to four inches up the grass stem or leaf.
- If you graze down to six to eight inches and move your sheep, you have evaded a lot of infective larvae. Let them die.
- Leaving behind sufficient residual also gives your grass a jump start in regrowth. You have left behind lots of photo-collectors (leaves). That is definitely not wasted grass!



Ready to move. Six to eight inches of residual. Photo: Pat Hansen

Ingestion: Leave Before the Larvae Arrive

It takes four to nine days after deposition for eggs in the fecal pellet to hatch into L3 larvae and then to crawl up on to their perch in the grass leaf.



L3 Infective larvae waiting for your sheep to ingest them. You can avoid them by exiting paddocks before L3 larvae get to their perch (four to nine days). Photo: Langston University Research and Extension

If you move animals from the paddock before four days, you miss the latest crop of L3 larvae. Simple as that.

One- to four-day paddock periods also give your irrigated (or rain-fed) grass production a huge boost. Try it and be amazed. Who doesn't want more carrying capacity?

Graze to Control: The Sum is Greater than the Parts

- Grazing antagonistically with the Barber Pole Worm's lifecycle yields big dividends. Use 40-day rest periods, long-length paddock residuals, and short grazing periods in combination to maximize effective grazing control. These same three practices will greatly improve the health of your soil and pastures. See ATTRA's *Building Healthy Pasture Soils*, <https://attra.ncat.org/attra-pub/summaries/summary.php?pub=580>.
- The incorporation of tannin-producing plants like birdsfoot trefoil in North-east pastures has been shown to reduce both fecal egg counts and FAMA-CHA scores, especially in conjunction with copper oxide wire particles (Parent et al., 2016). This may be worth investigating, especially in Utah, where birdsfoot trefoil grows relatively well. However, it is sometimes difficult to establish (Brummer et al., 2016).

- Multispecies grazing: In this situation, the Barber Pole Worm ends its life in a bovine's belly without shedding a single egg to your pasture. The same applies in reverse to bovine parasites (such as *Ostertagia*) that get vacuumed up by sheep. Can you employ this as a grazing strategy? See the ATTRA publication *Multi Species Grazing*, <https://attra.ncat.org/attra-pub/summaries/summary.php?pub=244>.
- Fence-line weaning is very effective in reducing lamb stress. Stress predisposes lambs to worm infection. This weaning strategy is not hard to do. Try it!



Fence-line weaning begun with ewes on the right and lambs on the left. Note that all heads are down, grazing. After 48 hours, the lambs were gently moved away. Not one bleat and little stress. Photo: Dave Scott, NCAT

Summary

Changing the way you graze can be one of the best moves you make to confront the Barber Pole Worm. To top it all off, there is little the worm can do genetically in your lifetime to adapt to it. You win.

References

Brummer, Joe, Jennifer MacAdam, Glen Shewmaker and M. Anowarul Islam. 2016. Establishing Birdsfoot Trefoil in the Mountain West. https://digitalcommons.usu.edu/cgi/viewcontent.cgi?referer=https://www.google.com/&httpsredir=1&article=2612&context=extension_curall

Parent, Molly, Tatiana Stanton, and Betsy Hodge. 2016. The Effect of Birdsfoot Trefoil Forage on Barber Pole Worm in Sheep and Goats in Northern NY. www.nnyagdev.org/wp-content/uploads/2011/07/BFTPrelim3doc.pdf

WormBoss. 2018. Factors Contributing to Paddock Contamination with Worms. www.wormboss.com.au/worms/roundworms/worms-on-pasture.php

ATTRA Resources

Managing Internal Parasites in Sheep and Goats
<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=215>

Multispecies Grazing
<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=244>

Tips for: Preventing Internal Parasites

<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=494>

Tools for Managing Internal Parasites in Small Ruminants: Pasture Management

<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=415>

Tools for Managing Internal Parasites in Small Ruminants: Copper Wire Particles

<https://attra.ncat.org/attra-pub/summaries/summary.php?pub=216>

Grazing to Control Parasites

By Dave Scott, NCAT Agriculture Specialist • Published June 2018. ©NCAT
IP565 • Slot 592



Produced by the National Center for Appropriate Technology
(parent organization to the ATTRA Project, www.attra.ncat.org)
www.ncat.org • 1-800-ASK-NCAT



This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2016-38640-25383 through the Western Sustainable Agriculture Research and Education program under subaward number EW 17-011. USDA is an equal opportunity employer and service provider.

Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture.



Contact Us! attra.ncat.org • Toll-free: 1-800-346-9140 • Email: askanag@ncat.org